



STOP THE SPREAD OF GERMS

Help prevent the spread of respiratory diseases like COVID-19.

Avoid close contact with people who are sick.



Cover your cough or sneeze with a tissue, then throw the tissue in the trash.



Avoid touching your eyes, nose, and mouth.



Clean and disinfect frequently touched objects and surfaces.



Stay home when you are sick, except to get medical care.



Wash your hands often with soap and water for at least 20 seconds.



For more information: www.cdc.gov/COVID19

8.25% Bleach Solutions

READ THE LABELS AND TAKE THE FOLLOWING STEPS TO ENSURE SAFETY IN YOUR CHILD CARE FACILITY

1	<p>Identify what bleach concentration is in your facility.</p> <p>Refer to the chart below for mixing instructions.</p> <p>Find the % sodium hypochlorite on the bottle.</p> <p>Avoid splash-less and scented bleaches.</p>	<p>Active Ingredient:</p> <p>Sodium Hypochlorite: 8.25%</p> <p>Other Ingredients: 91.75%</p> <p>Total: 100.00%</p>
----------	--	---

2	<p>Clean</p> <ul style="list-style-type: none"> Scrub with soap and warm water followed by a rinse with clean water. Always clean surfaces to remove visible soil, dirt, and contamination before using bleach solution. <i>Wipe dry with disposable towel.</i>
----------	---

3	<p>Mix</p> <ul style="list-style-type: none"> Mix fresh solutions daily for sanitizing and disinfecting. Add bleach to cool-water. Do not mix liquid bleach with other cleaning products, toilet bowl cleaners or ammonia, which may release hazardous gases in to the air.
----------	--

4	<p>Sanitize, Disinfect, Special Clean-up</p> <ul style="list-style-type: none"> Wet entire surface Leave solution on surface for 2 minutes for sanitizing and 5 minutes for disinfecting Dry with paper towel or allow to air-dry
----------	---

Sanitize (100 ppm)	Disinfect (600 ppm)	Special Clean-up (5000 ppm)
<p>CLEAN & SANITIZE AFTER EACH USE:</p> <ul style="list-style-type: none"> Mouthed toys and pacifiers Food service areas, dishes & utensils 	<p>CLEAN & DISINFECT AFTER EACH USE:</p> <ul style="list-style-type: none"> Diaper changing surface Toilet training seats 	<p>CLEAN & USE AS NEEDED FOR VOMIT AND DIARRHEA:</p> <p>Not for other bodily fluids</p>
<p>SANITIZE DAILY OR WHEN SOILED:</p> <ul style="list-style-type: none"> Dishcloths, synthetic sponges Food preparation appliances 	<p>DISINFECT DAILY OR WHEN SOILED:</p> <ul style="list-style-type: none"> Bathroom areas Common surfaces; floors, mats, tables, countertops and hard surface, door knobs, etc. 	<p>MIX SOLUTION WHEN NEEDED</p> <p>WEAR GLOVES AND MASKS TO PROTECT YOURSELF</p>
1/8 teaspoon bleach/ pint water	3/4 teaspoon bleach/ pint water	2 Tablespoons bleach/ pint water
1/4 teaspoon bleach/ quart water	1 1/2 teaspoon bleach/ quart water	4 Tablespoons bleach/ quart water
1 teaspoon bleach/ gallon water	2 Tablespoons bleach/ gallon water	1 cup (8 oz.) bleach/ gallon water



*For assistance with Sanitizing, Disinfecting, and Special Clean-up procedures, contact your regional Child Care Health Consultant through the Idaho Care Line by dialing 2-1-1 or 1-800-926-2588

*Caring for Our Children is available online at <http://efcc.nichdhs.org> or in print. This resource provides National health and safety recommendations. A routine cleaning schedule can be found in this resource as well.